arXiv:math/0510010v2 [math.DG] 11 Oct 2005

REDUCTION OF TWISTED GENERALIZED KÄHLER STRUCTURE

YI LIN, SUSAN TOLMAN

ABSTRACT. We show that our earlier work in [LT05] extends to the twisted case, that is, we defined a notion of moment map and reduction in both twisted generalized complex geometry and twisted generalized Kähler geometry.

This is a short note to show that our results in our previously posted paper, *Symmetries in generalized Kähler geometry* [LT05], can be easily extended to the case of twisted generalized geometries with only minor modifications. In particular, we define twisted generalized complex reduction and twisted generalized Kähler reduction. We hope that this goes some way towards providing the framework which Kapustin and Li suggested would be useful [KL].

This note is not intended as a completely separate paper, but should rather be read in conjunction with [LT05]. In particular, we will not repeat the brief history of this subject which can be found there, except to reiterate that it was first introduced in [H02] and developed much further in [Gua04].

However, it is important to mention here that closely related works which have appeared since we first posted. We have not had an opportunity to carefully read these papers but would like to at least attempt to describe some basic similarities and differences.

Shengda Hu wrote a related paper [Hu05], which was partially inspired by an early version of our paper [LT05] which we gave him in early June. His paper includes a notion of twisted complex reduction which is very similar to ours in the untwisted case. More generally, he considers twisted complex structures in the framework of Hamiltonian symmetry. The work in the current note was completed after his paper appeared, so also cannot be considered as fully independent. Our main motivation is to demonstrate twisted generalized Kähler reduction, which is not in [Hu05]. However, it is worth noting that even in the twisted generalized complex case our results are slightly different.

In [SX05], Stiénon and Ping independently develop notions of generalized complex and Kähler reduction which seems rather different from ours in both perspective and techniques. In particular, instead of working with generalized moment maps, they consider quotients of arbitrary subsets; so

---

Date: February 2, 2008.

our theorems do not appear in their paper (or vise-versa). They do not consider bi-Hermitian or twisted structures.

In [BCG05], which is also independent of our work, Busztychn, Cavalcanti, and Gualtieri define a quite general notion of the reduction of Courant algebroids and Dirac structures which includes twisted generalized complex and twisted generalized Kähler reduction as special cases. They work with both arbitrary subsets and moment maps, but again their techniques are different from ours. They also construct two bi-Hermitian structures on $\mathbb{CP}^2$, but do not construct bi-Hermitian structures on the other spaces we consider in [LT05].

We begin with a brief introduction, following [Gua04]. (We would like to mention that in addition to the many concrete instances where we quote Gualtieri, our whole perspective on generalized geometries was heavily influenced by his excellent thesis on this subject.)

Let M be a n dimensional manifold. There is a natural metric of type (n, n) on $TM \oplus T^*M$ given by

$$\langle X + \alpha, Y + \beta \rangle = \frac{1}{2}(\alpha(Y) + \beta(Y)),$$

which extends naturally to $T_{\mathbb{C}}M \oplus T^*_{\mathbb{C}}M = (TM \oplus T^*M) \otimes \mathbb{C}$.

A **generalized almost complex structure** on a manifold M is a bundle map $\mathcal{J}\colon TM \oplus T^*M \to TM \oplus T^*M$ which is orthogonal with respect to the natural metric defined above so that $\mathcal{J}^2 = -1$.

For a closed form $H \in \Omega^3(M)$, the **H-twisted Courant bracket** on $T_{\mathbb{C}}M \oplus T^*_{\mathbb{C}}M$ is given by

$$[X + \alpha, Y + \beta]_H = [X, Y] + L_X\beta - L_Y\alpha - \frac{1}{2}(d\iota_X\beta - d\iota_Y\alpha) + \iota_Y\iota_X H.$$

A **H-twisted generalized complex structure** on M is a generalized almost complex structure $\mathcal{J}$ so that the $\sqrt{-1}$ eigenbundle of $\mathcal{J}$ is closed under the H-twisted Courant bracket. And a **H-twisted generalized Kähler structure** on a manifold M is a pair of commuting **H**-twisted generalized complex structures $\mathcal{J}_1$ and $\mathcal{J}_2$ on M so that $-\mathcal{J}_1\mathcal{J}_2$ is a positive definite bundle map on $TM \oplus T^*M$ with respect to the natural metric.

As the following example shows, the concept of twisting is most interesting when H represents a non-trivial cohomology class.

Example 1. [Gua04] Given a two-form B on a manifold M, consider the orthogonal bundle map $TM \oplus T^*M \to TM \oplus T^*M$ defined by

$$e^B = \begin{pmatrix} 1 & 0 \\ B & 1 \end{pmatrix},$$

where B is regarded as a skew-symmetric map from TM to T^*M. If $\mathcal{J}$ is a generalized almost complex structure on M, then $\mathcal{J}' = e^B\mathcal{J}e^{-B}$ is another generalized almost complex structure on M, called the **B-transform** of $\mathcal{J}$. For any $B \in \Omega^2(M)$ and closed $H \in \Omega^3(M)$, the B-transform of a H-twisted generalized complex (or Kähler) structure is a $H + dB$-twisted generalized complex (or Kähler) structure.

It will also be convenient to have the following definition.

Definition 2. *Let a compact Lie group* G *act on a manifold* M*. The* **Cartan model** *for the equivariant cohomology of* M *is defined as follows: The degree* n *co-chains are*

$$\Omega^n_G(M) = \bigoplus_i (\Omega^{n-2i}(M) \otimes S^i(\mathfrak{g}^*))^G,$$

where S^i *denotes polynomials of degree* i*. The differential* $d_G : \Omega^n_G \to \Omega^{n+1}_G$ *is defined by*

$$d_G(\alpha \otimes p)(\xi) = (d\alpha - \iota_{\xi_M}\alpha)p(\xi) \qquad \text{for all } \xi \in \mathfrak{g},$$

where we think of $\Omega^*_G(M)$ *as the space of equivariant polynomial mappings from* $\mathfrak{g}$ *to* $\Omega^*(M)$*. (If* G *acts on a vector space* A*, let* A^G *denote the invariant subspace.) The* **equivariant cohomology** *of* M *is* $H^*_G(M) = H^*(\Omega^*_G, d_G)$*.*

Remark 3. Let a compact Lie group G act on a manifold M so that it acts freely on a submanifold $L \subset M$. There is a natural map, called the **Kirwan map**

$$\kappa : H^*_G(M) \to H^*(L/G)$$

which is the composition of the restriction map from $H^*_G(M)$ to $H^*_G(L)$ with the natural isomorphism from $H^*_G(L)$ to $H^*(L/G)$.

A form $B \in \Omega^n(M)$ is **basic** if it is invariant and $\iota_{\xi_M} B = 0$ for all $\xi \in \mathfrak{g}$. Then B **descends** to $\widetilde{B} \in \Omega^n(L/G)$, that is, the pull-back of $\widetilde{B}$ to L is the restriction of B.

If $B \in \Omega^n(M)^G \subset \Omega^n_G(M)$ is **equivariantly closed**, that is, $d_G B = 0$, then B is closed and basic and $\kappa[B] = [\widetilde{B}]$. More generally, if $\eta \in \Omega^n_G(M)$ is equivariantly closed, there exists $\Gamma \in \Omega^{n-1}_G(L)$ so that $\eta|_L + d_G\Gamma \in \Omega^n(L)^G \subset \Omega^n_G(L)$. Since $\eta + d_G\Gamma$ is equivariantly closed, it descends to $\widetilde{\eta} \in \Omega^n(L/G)$ and $\kappa[\eta] = [\widetilde{\eta}]$.

It is easy to check that Lemma 2.1 and Lemma 2.2 in [LT05] still hold for the twisted Courant bracket, and that the same proofs work. More specifically, if $f : M \to g^*$ is a submersion, then $df^{\perp}_{\mathbb{C}} \subset T_{\mathbb{C}}M \oplus T^*_{\mathbb{C}}M$ is closed under the H-twisted Courant bracket for any closed $H \in \Omega^3(M)$. Therefore, if Γ is a sub-bundle of $T_{\mathbb{C}}M \oplus T^*_{\mathbb{C}}M$ which is closed under the H-twisted Courant bracket, then the image of $\Gamma \cap df^{\perp}_{\mathbb{C}}$ in $T_{\mathbb{C}}(f^{-1}(0)) \oplus T^*_{\mathbb{C}}(f^{-1}(0))$ is closed under the $H|_{f^{-1}(0)}$-twisted Courant bracket. Similarly, let G act freely on M and assume that $H \in \Omega^3(M)$ is closed and basic. Then H descends to a closed form $\widetilde{H} \in \Omega^3(M/G)$ and the set of G-invariant sections of $(\mathfrak{g}_M)^{\perp}_{\mathbb{C}}$ is closed under the H-twisted Courant bracket. Therefore, if Γ is a sub-bundle of $T_{\mathbb{C}}M \oplus T^*_{\mathbb{C}}M$ which is closed under the H-twisted Courant bracket, then the image of $\Gamma \cap (\mathfrak{g}_M)^{\perp}_{\mathbb{C}}$ in $T_{\mathbb{C}}(M/G) \oplus T^*_{\mathbb{C}}(M/G)$ is closed under the $\widetilde{H}$-twisted Courant bracket.

In this context, we will work with a variant of the notion of generalized moment map we defined in [LT05].

Definition 4. *Let a compact Lie group* G *with Lie algebra* $\mathfrak{g}$ *act on a manifold* M*, preserving an* H*-twisted generalized complex structure* $\mathcal{J}$*, where* $H \in \Omega^3(M)^G$ *is closed. Let* $L \subset T_{\mathbb{C}}M \oplus T^*_{\mathbb{C}}M$ *denote the* $\sqrt{-1}$ *eigenbundle of* $\mathcal{J}$*. A* **twisted generalized moment map** *is a smooth function* $f\colon M \to \mathfrak{g}^*$ *so that*

- *There exists a one form* $\alpha \in \Omega^1(M, \mathfrak{g}^*)$*, called the* **moment one form***, so that* $\xi_M - \sqrt{-1}\,(df^\xi + \sqrt{-1}\,\alpha^\xi)$ *lies in* L *for all* $\xi \in \mathfrak{g}$*, where* ξ_M *denote the induced vector field.*
- f *is equivariant.*

Example 5.

a) Let G act on a generalized complex manifold with generalized moment map $f + \sqrt{-1}\,h$. Then f is a twisted generalized moment map with moment one form dh.
b) Let G act on an H-twisted generalized complex manifold $(M, \mathcal{J})$ with twisted generalized moment map f and moment one-form α. If $B \in \Omega^2(M)^G$, then G acts on the B-transform of $\mathcal{J}$ with twisted generalized moment map f and moment one form α', where $(\alpha')^\xi = \alpha^\xi + \iota_{\xi_M} B$ for all $\xi \in \mathfrak{g}$.

Let a compact Lie group G act on a twisted generalized complex manifold $(M, \mathcal{J})$ with twisted generalized moment map f. Let $\mathcal{O}_a$ be the co-adjoint orbit through $a \in \mathfrak{g}^*$. If G acts freely on $f^{-1}(\mathcal{O}_a)$, then $\mathcal{O}_a$ consists of regular values and $M_a = f^{-1}(\mathcal{O}_a)/G$ is a manifold, which we still call the **generalized complex quotient**. The proof of Lemma 3.8 in [LT05] applies almost word for word to the following generalization.

Lemma 6. *Let a compact Lie group* G *act on an* H*-twisted generalized complex manifold* $(M, \mathcal{J})$ *with a twisted generalized moment map* $f\colon M \to \mathfrak{g}^*$*. Let* $\mathcal{O}_a$ *be a co-adjoint orbit through* $a \in \mathfrak{g}^*$ *so that* G *acts freely on* $f^{-1}(\mathcal{O}_a)$*. Assume that the moment one-form is trivial and* H *is basic. Then* H *descends to* $\widetilde{H} \in \Omega^3_0(M_a)$ *and the generalized complex quotient* M_a *naturally inherits a* $\widetilde{H}$*-twisted generalized complex structure* $\widetilde{\mathcal{J}}$*.*

Moreover, for all $m \in f^{-1}(\mathcal{O}_a)$*,*

$$\operatorname{type}(\widetilde{\mathcal{J}})_{[m]} = \operatorname{type}(\mathcal{J})_m.$$

Example 7.

a) Let a compact Lie group act on a (untwisted) generalized complex manifold with a real generalized moment map. Then Lemma 3.8 in [LT05] and Lemma 6 yield identical generalized complex structures on M_a.
b) Let a compact Lie group G act on an H-twisted generalized complex manifold $(M, \mathcal{J})$ with twisted generalized moment map $f : M \to \mathfrak{g}^*$. Assume that the moment one-form is trivial and H is

basic. If $B \in \Omega^2(M)$ is basic, then it descends to $\widetilde{B} \in \Omega^2(M_a)$, the B-transform of $\mathcal{J}$ satisfies the same conditions, and its twisted generalized complex quotient is the $\widetilde{B}$-transform of the twisted generalized complex quotient of $\mathcal{J}$.

Lemma 8. *Let a compact Lie group* G *act freely on a manifold* M. *Let* H *be an invariant closed three form and let* α *be an equivariant mapping from* $\mathfrak{g}$ *to* $\Omega^1(M)$. *Fix a connection* $\theta \in \Omega(M, \mathfrak{g})$. *Then if* $H + \alpha \in \Omega^3_G(M)$ *is equivariantly closed, there exists a natural form* $\Gamma \in \Omega^2(M)^G$ *so that* $\iota_\xi \Gamma = \alpha^\xi$. *Thus* $H + \alpha + d_G\Gamma \in \Omega^3(M)^G \subset \Omega^3_G(M)$ *is closed and basic and so descends to a closed form* $\widetilde{H} \in \Omega^3(M/G)$ *so that* $[\widetilde{H}]$ *is the image of* $[H + \alpha]$ *under the Kirwan map.*[1]

Proof. Since $H + \alpha$ is equivariantly closed

- $dH = 0$.
- $\iota_{\xi_M} H = d\alpha^\xi$ for all $\xi \in \mathfrak{g}$.
- $\iota_{\xi_M} \alpha^\eta = -\iota_{\eta_M} \alpha^\xi$ for all ξ and η in $\mathfrak{g}$.

Define $\beta \in \Omega^2(M)^G$ by $\beta(X, Y) = -\alpha^{\theta(X)}(\theta(Y)_M) = \alpha^{\theta(Y)}(\theta(X)_M)$ for every vector field X and Y. Define $\Gamma = \Gamma_\theta = -(\alpha, \theta) + \beta \in \Omega^2(M)^G$. Then $\iota_{\xi_M}\Gamma = \alpha^\xi$ for all $\xi \in \mathfrak{g}$. Since Γ is G-invariant, $\iota_{\xi_M} d\Gamma = -d\iota_{\xi_M}\Gamma = -d\alpha^\xi = -\iota_{\xi_M} H$. Therefore, $H + d\Gamma$ is basic. □

We can now prove the twisted version of Proposition 3.10.

Proposition 9. *Let a compact Lie group* G *act on an* H*-twisted generalized complex manifold* $(M, \mathcal{J})$ *with twisted generalized moment map* $f : M \to \mathfrak{g}^*$ *and moment one-form* $\alpha \in \Omega^1(M, \mathfrak{g}^*)$. *Let* $\mathcal{O}_a$ *be a co-adjoint orbit through* $a \in \mathfrak{g}^*$ *so that* G *acts freely on* $f^{-1}(\mathcal{O}_a)$. *Assume that* $H + \alpha$ *is equivariantly closed. Given a connection on* $f^{-1}(\mathcal{O}_a)$, *the twisted generalized complex quotient* M_a *inherits a* $\widetilde{H}$*-twisted generalized complex structure* $\widetilde{\mathcal{J}}$, *where* $\widetilde{H}$ *is defined as in the Lemma above. Up to* B*-transform,* $\widetilde{J}$ *is independent of the choice of connection. Finally, for all* $m \in f^{-1}(\mathcal{O}_a)$,

$$\mathrm{type}(\widetilde{\mathcal{J}})_{[m]} = \mathrm{type}(\mathcal{J})_m.$$

Proof. By restricting to a neighborhood of $f^{-1}(\mathcal{O}_a)$, we may assume that G acts freely on M. Given a connection $\theta \in \Omega^1(M, \mathfrak{g})$, by the lemma above there exists $\Gamma \in \Omega^2(M)$ so that $\iota_{\xi_M}\Gamma = \alpha^\xi$. So the Γ-transform of $\mathcal{J}$ is a $H - d\Gamma$-twisted generalized complex structure with twisted generalized moment map f and trivial moment one-form, and hence induces a natural $\widetilde{H}$-twisted generalized complex structure $\widetilde{\mathcal{J}}$ on M_a by Lemma 6.

Given another connection θ', then $\Gamma_\theta - \Gamma_{\theta'}$ is basic and hence descends to $\widetilde{\gamma} \in \Omega^2(M)$. Consequently, the resulting twisted generalized complex structure on M_a is the $\widetilde{\gamma}$ transform of $\widetilde{\mathcal{J}}$. □

---

[1] In fact, the analogous statement holds for any equivariantly closed form, but the proof in general is more involved.

Remark 10. In particular, even if $H = 0$, if $[\alpha] \neq 0$ then in general $[\widetilde{H}]$ will also not vanish. Thus, in principle it may be possible to get non-trivially twisted quotients from non-twisted spaces.

Example 11.

a) Let a compact Lie group act on a generalized complex manifold with generalized moment map $f + \sqrt{-1}\, h$. The generalized complex structure on M_a induced by Proposition 9 is the $(h, d\theta)$-transform of the generalized complex structure induced by Proposition 9. In particular, it may be twisted.

b) Let G act on an H-twisted generalized complex manifold $(M, \mathcal{J})$ with twisted generalized moment map f and moment one-form α. Assume that $H + \alpha$ is equivariantly closed. Fix a connection θ on $f^{-1}(\mathcal{O}_a)$. Given any $B \in \Omega^2(M)^G$, let $\Gamma \in \Omega^2(M)^G$ be the natural form associated to the closed form $d_G(B) \in \Omega^3_G(M)$ by Lemma 8 . Then $B + \Gamma$ is basic and hence descends to a form $\widetilde{B} \in \Omega^2(M_a)$, and the twisted generalized complex quotient of the B-transform of $\mathcal{J}$ is the $\widetilde{B}$ transform of the twisted generalized complex quotient. Hence, we cannot get interesting new examples by applying B-transforms to the space upstairs.

A **twisted generalized moment map** and **moment one-form** for a group action on a twisted generalized Kähler manifold $(M, \mathcal{J}_1, \mathcal{J}_2)$ are simply a twisted generalized moment map and moment one-form for the twisted generalized complex structure $\mathcal{J}_1$. We are now ready to prove our final proposition.

Proposition 12. *Let a compact connected Lie group* G *act on an* H*-twisted generalized Kähler manifold* $(M, \mathcal{J}_1, \mathcal{J}_2)$ *with twisted generalized moment map* $f\colon M \longrightarrow \mathfrak{g}^*$ *and moment one-form* $\alpha \in \Omega^1(M, \mathfrak{g}^*)$. *Let* $\mathcal{O}_a$ *be a co-adjoint orbit through* $a \in \mathfrak{g}^*$ *so that* G *acts freely on* $f^{-1}(\mathcal{O}_a)$. *Assume* $H+\alpha \in \Omega^3_G(M)$ *is equivariantly closed. Then the generalized Kähler quotient* M_a *naturally inherits a* $\widetilde{H}$*-twisted generalized Kähler structure* $(\widetilde{\mathcal{J}_1}, \widetilde{\mathcal{J}_2})$, *where* $\widetilde{H}$ *is defined as in Lemma 8 using the canonical connection on* M.

Finally, let $\mathfrak{k}$ *be the Lie algebra of the stabilizer* K *of* a, *and let* L_2 *be the* $\sqrt{-1}$ *eigenbundle of* $\mathcal{J}_2$. *Then for all* $m \in M$,

$$\operatorname{type}(\widetilde{\mathcal{J}_1})_{[m]} = \operatorname{type}(\mathcal{J}_1)_m, \qquad \textit{and}$$

$$\operatorname{type}(\widetilde{\mathcal{J}_2})_{[m]} = \operatorname{type}(\mathcal{J}_2)_m - \frac{1}{2}\dim G - \frac{1}{2}\dim(K) + 2\dim(\mathfrak{h}_M \cap \pi(L_2))_m.$$

Proof. By restricting to a neighborhood of $f^{-1}(\mathcal{O}_a)$, we may assume that G acts freely on M. The Γ_θ transform of $(\mathcal{J}_1, \mathcal{J}_2)$ is a $H + d\Gamma_\theta$-twisted generalized Kähler structure with twisted generalized moment map f and trivial moment one-form. From this point, the proof of Proposition 4.6 in [LT05] goes through without change. □

REFERENCES

[BCG05] H. Bursztyn, G. Cavalcanti, and M. Gualtieri, *Reduction of Courant algebroids and generalized complex structures*, math.DF

[Cr04] M. Crainic, *Generalized complex structures and Lie brackets*, math.DG/04112097

[Gua04] M. Gualtieri, *Generalized complex geometry*, math.DG/0411221

[LT05] Y. Lin, S. Tolman *Symmetries in generalized Kähler geometry*, math/0509069

[H02] N. Hitchin, *Generalized Calabi-Yau manifolds*, math.DG/0209099

[H05] N. Hitchin, *Instantons, Poisson structures and generalized Kahler geometry*, math.DG/0503432

[Hu05] S. Hu, *Hamiltonian symmetries and reduction in generalized geometry*, math.DG/0509060

[KL] A. Kapustin and Y. Li, *Topological sigma-models with H-flux and twisted generalized complex manifolds*, hep-th/0407249

[SX05] M. Stiénon and X. Ping, *Reduction of generalized complex structure*, math.DG/0509393

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, CANADA, M5S2E4, YILIN@MATH.TORONTO.EDU

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, URBANA, IL, USA, 61801, STOLMAN@MATH.UIUC.EDU